



BC Storm Surge Forecasting System

2020-21 Storm Surge Almanac

Climate outlook and tidal elevations for fall/winter 2020

October 16, 2020

BC Storm Surge Forecasting System

Executive Summary

The 2019-20 storm surge season was marked by several weak to moderate wind events spaced sporadically through the winter. Early 2020 saw the strongest wind events before cooler higher-pressure conditions set in. The weather last year was influenced by ENSO-neutral conditions in the Pacific basin. **The 2020-21 season is under a La Niña advisory expected to persist through the winter of 2020.**

The strongest wind storm on January 16th brought a peak surge of only 58 cm which ranked 46th of the last 57 years with a 1.3 year return period (Table 1). **The current La Niña conditions are normally associated with a weakened jet stream and a storm track that is displaced north of its normal winter location.** However, this year has already seen several strong, closed-low pressure systems hit the southern BC coast, which may be an indication of an active late fall/early winter storm season.

The Pacific Decadal Oscillation is now turning negative after being predominantly positive since 2014. Current coastal sea surface height (SSH) anomalies are near zero at the coast, but somewhat elevated well offshore related to the persistent anomalously warm surface waters in the north Pacific.

Annual coastal flooding risk is greatest during the seasonal perigean spring tides, which generate extreme high tidal levels during the winter months. This winter (2020-21) the highest tides for the Lower Mainland are concentrated in the third weeks of November, December and January. For Victoria, the highest tides occur mid-November, mid- and late-December and mid-January.

BC Storm Surge Forecasting System

Table 1. Rank of annual maximum storm surge at Point Atkinson (out of the past 57 years) for years since 1997. Reference year represents the year at the start of the winter season.

Year	Rank (out of last 56 years)	Surge (m)	ENSO Phase (red El Nino, blue La Nina)
1998	1	1.03	moderate-strong
2015	3	0.95	very strong
2006	5	0.91	weak
2002	8	0.89	moderate
2016	9	0.85	moderate-strong
2011	11	0.84	weak
2005	14	0.82	weak
2018	16	0.81	moderate
2007	21	0.79	weak-moderate
2014	24	0.77	neutral
2001	27	0.74	neutral
2009	31	0.72	moderate
2012	35	0.68	neutral
1997	37	0.66	very strong*
2010	40	0.65	moderate-strong
2019	46	0.58	neutral
2004	47	0.57	weak
2003	50	0.57	neutral
2017	51	0.55	neutral
2013	52	0.55	weak
1999	54	0.53	strong
2000	55	0.46	weak
2008	57	0.44	moderate

BC Storm Surge Forecasting System

Overview

This document provides a summary of the current and expected meteorological and oceanographic conditions for southern British Columbia as they relate to the 2020-21 winter storm season. The focus is on total water level (tide plus surge), and the risk of total water level approaching or exceeding the historical maximum observed values at Point Atkinson (Vancouver) and Victoria.

Fall weather has been intermittently wet and dry accompanied by several fast moving compact wind storms. The summer was dry but fall rains have returned water levels in most rivers and reservoirs to near-normal flows and volumes. Storm surge can be a confounding factor during many late-fall and winter storms when streams are full and hydro reservoir levels are near capacity. The additional barrier of high tide and surge at the mouth of rivers can adversely influence the ability to shed water at the time when discharge is most required.

The main factors that contribute to extreme marine coastal water levels are seasonal high tides (“king tides”), storm intensity (high wind velocity and low atmospheric pressure) and positive coastal sea level anomalies (departures from the long-term mean) due to basin-scale ocean climate phenomena such as the El Niño/Southern Oscillation (ENSO). Each of these risk factors is discussed in the following sections.

BC Storm Surge Forecasting System

Climatological Factors

Large-scale climatological phenomena affect both the weather and climate. The statistics of synoptic scale meteorological conditions in a given year may be influenced by the larger scale atmospheric and oceanographic conditions that are often described in terms of select climate indices. Two indices of significance to the west coast of North America are the Pacific Decadal Oscillation (PDO) and the El Niño/Southern Oscillation (ENSO)(Table 2). A description of these phenomena and a discussion of their potential relationship to the winter storm surge statistics for British Columbia are included in the Appendix.

Table 2. Monthly Pacific Decadal Oscillation (PDO) and multi-variate ENSO index values from 2016-present.

Year	2016		2017		2018		2019		2020	
	PDO	ENSO	PDO	ENSO	PDO	ENSO	PDO	ENSO	PDO	ENSO
Jan	1.53	2.23	0.77	-0.06	0.70	-0.62	0.45	0.08	-0.19	0.30
Feb	1.75	2.17	0.70	-0.06	0.37	-0.73	0.16	0.52	-0.41	0.30
Mar	2.40	1.96	0.74	-0.08	-0.05	-0.50	0.30	0.77	-0.86	0.2
Apr	2.62	2.07	1.12	0.77	0.11	-0.43	0.76	0.33	-0.62	-0.1
May	2.35	1.70	0.88	1.46	0.11	0.47	0.97	0.26	0.25	-0.2
Jun	2.03	1.00	0.79	1.05	-0.04	0.47	0.95	0.35	0.14	-0.7
Jul	1.25	0.31	0.10	0.46	0.11	0.08	1.28	0.24	-0.17	-1
Aug	0.52	0.18	0.09	0.03	0.18	0.13	0.92	0.30	-0.44	-1
Sep	0.45	-0.10	0.32	-0.45	0.09	0.51	1.03	0.20	-0.14	-1.2
Oct	0.56	-0.38	0.05	-0.55	0.18	0.41	-0.22	0.30		
Nov	1.88	-0.21	0.15	-0.28	0.13	0.26	0.44	0.50		
Dec	1.17	-0.12	0.50	-0.58	0.39	0.13	0.75	0.40		

Pacific Decadal Oscillation

In 2000, the PDO shifted away from the multi-decadal warm phase (1977-2000), to what had been a predominantly a cold phase (negative PDO) until 2014. Strongly positive PDO values recorded between January 2014 and July 2016 due primarily to El Niño events have largely remained positive since 2017. **The PDO index as of September 2020 is -0.14.**

El Niño/Southern Oscillation

The NCEP Climate Prediction Center (CPC) provides regular updates of the state of ENSO in the Pacific Ocean. The October 8, 2020 bulletin from the ENSO alert system indicates a La Niña advisory. **The current ENSO MEI level (as of September 2020) is -1.2.** Consensus model forecasts published by CPC (Fig. 1) suggest an 85% chance of La Niña conditions through the 2020 fall/winter, and a 60% chance of La Niña conditions continuing through to the Northern Hemisphere spring. The complete commentary is available at:

BC Storm Surge Forecasting System

http://www.cpc.ncep.noaa.gov/products/analysis_monitoring/enso_advisory/ensodisc.pdf

From the CPC commentary of October 8, 2020 (Figure references in the quoted text are for the commentary at the link above):

“La Niña continued during September, as evidenced by below-average sea surface temperatures (SSTs) extending from the Date Line to the eastern Pacific Ocean... The atmospheric circulation anomalies over the tropical Pacific Ocean remained consistent with La Niña. Low-level wind anomalies were easterly across most of the tropical Pacific, and upper-level wind anomalies were westerly over the east-central Pacific. ... Also, both the Southern Oscillation and Equatorial Southern Oscillation indices remained positive. Overall, the coupled ocean-atmosphere system indicates the continuation of La Niña.

A majority of the models in the IRI/CPC plume predict La Nina (Niño-3.4 index less than -0.5°C) to persist through the Northern Hemisphere winter 2020-21 and to weaken during the spring [Fig. 6]. The latest forecasts from several models, including the NCEP CFSv2, suggest the likelihood of a moderate or even strong La Nina (Niño-3.4 index values $< -1.0^{\circ}\text{C}$) during the peak November-January season. The forecaster consensus supports that view in light of significant atmosphere-ocean coupling already in place. In summary, La Niña is likely to continue through the Northern Hemisphere winter 2020-21 [...].”




Figure 6. Forecasts of sea surface temperature (SST) anomalies for the Niño 3.4 region (5°N - 5°S , 120°W - 170°W). Figure updated 18 September 2020.

Figure 1: [CPC Figure 6] CPC numerical model consensus forecast for ENSO conditions for Fall 2020 to Spring 2021. The consensus is for an 85% chance of ENSO-neutral conditions through winter 2020.

BC Storm Surge Forecasting System

Eichler and Higgens (2006 - referred to hereafter as EH2006) compared North American extratropical storm activity to ENSO phases based on the NCEP reanalysis period of 1950-2002 and the European ECMWF 40-year reanalysis data from 1971-2000. They categorized years based on an ENSO Intensity Scale (EIS) equal to twice the “Oceanic Niño Index” (see Table 3).

Table 3: (From Eichler and Higgens (2006) Table 1)

EIS > 3	strong El Niño	1958, 1966, 1973, 1983, 1992, 1998, 2003
0 < EIS < 3	weak/moderate El Niño	1964, 1969, 1977, 1978, 1987, 1988, 1995
EIS = 0	Neutral	1952, 1953, 1954, 1959, 1960, 1961, 1963, 1967, 1970, 1979, 1980, 1981, 1982, 1986, 1990, 1991, 1993, 1994, 1997, 2002
-3 < EIS < 0	weak/moderate La Niña	1951, 1955, 1957, 1962, 1965, 1968, 1971, 1972, 1975, 1984, 1985, 1996, 2001
EIS < -3	strong La Niña	1950, 1956, 1974, 1976, 1989, 1999, 2000

Over the entire reanalysis period, EH2006 provide an average seasonal frequency of the number of storms in the northeast Pacific that impact the southern coast of British Columbia. Results show an average of three storms in the period spanning October to December (OND) and four in the period from January to March (JFM). When the storms were binned by ENSO phase (as described in Table 3), the JFM storm track frequency was shown to be highest (four or more) during a moderate or strong La Niña phase and during a strong El Niño phase. The lowest storm frequency in the region (4 or less) occurs, on average, during ENSO neutral or moderate El Niño conditions. This analysis was not done for the OND period so it is not clear from the discussion of EH2006 whether or not the storms have shifted earlier in the season, but it is assumed that the weakening of storm frequency in JFM during ENSO neutral to moderate El Niño occurs in all seasons. **Winter forecast for 2020-21 is for a moderate La Niña.**

BC Storm Surge Forecasting System

Surge Statistics

Residual sea level (water level - tidal height) statistics for southern British Columbia are examined for Point Atkinson and Victoria from 1980 to present. This time period was chosen because it represents a modern era in the gauge data. One drawback of using this period is that the beginning coincides with the start of the 23-year PDO warm phase (1977-2000), and therefore may include climatological biases (weighted towards warm PDO conditions). The tides were calculated for both tide gauge stations and subtracted from observed water levels. The resulting water level residuals were binned in 5 cm bins; all residuals greater than 20 cm were retained for the analysis.

Point Atkinson

The maximum residual water level measured (1980-present) at Pt. Atkinson is **1.03 m in March, 1999**. For residuals > 20 cm, the 90th percentile corresponds to residuals >45 cm. The percentage and cumulative percentage represented for each 5-cm bin are shown in Fig. 2.




Figure 2. Residual water levels at Pt. Atkinson (>20 cm) in 5-cm bins as a percentage of the total number of residuals >20 cm (blue) based on hourly observations. The cumulative percentage at each bin level is shown in orange. Residuals >45 cm lie within the 90th percentile (black line) of all residuals >20 cm.

BC Storm Surge Forecasting System

Victoria

The maximum residual water level measured (1980-present) at Victoria is **0.80 m in January 1983**. The 90th percentile comprises all residuals >40 cm. The percentage and cumulative percentage represented for each 5-cm bin are shown in Fig. 3.




Figure 3. Residual water levels at Victoria (>20 cm) in 5-cm bins as a percentage of the total number of residuals >20 cm (blue) based on hourly observations. The cumulative percentage at each bin level is shown in orange. Residuals >40 cm lie within the 90th percentile (black line) of all residuals >20 cm.

BC Storm Surge Forecasting System

Tides

The highest tides of the year, known as the perigean spring tides, occur near the summer and winter solstices. High tides during this period reach (or slightly exceed) 5.0 m at Point Atkinson and 3.1 m at Victoria respectively. Water levels of this magnitude leave coastal areas vulnerable to flooding by storm surge. To illustrate the importance of these periods of extreme tidal height, a storm surge of 0.6 m (1.3 year return period) concurrent with peak winter tides (5.0 m) would cause a total water level equivalent to the 1982 record high at Point Atkinson (and the 2003 record high at Victoria), even without the presence of El Niño-induced elevated coastal sea levels.

Point Atkinson Tide Calendar

The historical recorded highest water level at Point Atkinson occurred on December 16, 1982 when the water gauge measured a total water level of 5.61 m (4.71 m tide + 0.90 m anomaly). The factors that led to the historical highest water level were a combination of high seasonal tide, strong winds, low atmospheric pressure and a coastal sea-level height anomaly of approximately 0.2 m driven by one of the most intense El Niño events on record.

Figure 4 highlights the dates when peak tides exceed 4.7 m relative to station datum at Point Atkinson; these periods offer broad windows during which a moderate or high storm surge could result in extreme water levels. Days with predicted tides in the range of 4.7-4.8 m are shaded yellow, while those with tides in the range of >4.9 m above chart datum are shaded red.

The periods where predicted tide exceeds 4.9 m at Point Atkinson occur from November 17-19, December 15-19 and January 12-16.

BC Storm Surge Forecasting System




Figure 4. 2020-21 calendar dates for tides exceeding 4.7 m at Point Atkinson.

Victoria Tide Calendar

The historical maximum observed water level at Victoria of 3.71 m above chart datum (3.14 m tide + 0.57 m surge) occurred on January 2, 2003. This coincided with the time of highest seasonal tide and during a moderate El Niño.

Figure 5 highlights the dates when peak tides exceed 2.8 m relative to station datum (dates where tides >3.0 m are shaded red). Depending on the predominant wind direction, several regions around Victoria are at risk to storm surge damage including Fairfield (Ross Bay), Oak Bay and Cadboro Bay. This year, **the highest predicted tides (>3.0 m) occur during December 14-17, December 12-14, December 29-31, January 8-11 and February 8.**

BC Storm Surge Forecasting System




Figure 5. 2020-21 calendar dates for tides exceeding 2.8 m at Victoria.

BC Storm Surge Forecasting System

Sea Surface Height Anomaly

Sea surface height (SSH) anomalies derived from the JASON radar altimetry satellite are processed by the Jet Propulsion Laboratory (JPL) at the California Institute of Technology. The anomalies show the interannual variability of SSH after the mean and seasonal signals and the trend have been removed. The SSH anomalies are used to highlight large scale spatial trends, often caused by climatological phenomena such as ENSO.

The ENSO-neutral conditions of 2019-20 have shifted to moderate La Niña conditions. Near equatorial SSH anomalies are negative, and forecasts are for continuing La Niña conditions through spring 2021. Currently, SSH anomalies are near zero all along the North American west coast (Fig. 6), but a strongly positive anomaly (10-15 cm) related to steric effects from surface warming remains in the north Pacific.




Figure 6. JPL image of 10-day averaged SSH anomalies over the Pacific Ocean on September 22, 2020. Equatorial SSH Anomalies are consistent with La Niña conditions.

The latest seasonal (July-October) SST anomaly (Fig. 7) published by the Earth System Research Laboratory (ERSL/NOAA) shows residual warming throughout the northeast Pacific and equatorial SST anomalies consistent with La Niña.

BC Storm Surge Forecasting System




Figure 7. ERSL/NOAA seasonal SST anomaly for July 12-October 10, 2020 (<https://psl.noaa.gov/map/clim/sst.shtml>).

BC Storm Surge Forecasting System

Summary

From the information available as of October 2020, the winter 2020-21 storm surge season will be characterized by the following:

1. **ENSO:** La Niña conditions to continue through to spring.
2. **PDO:** slightly negative and trending downward
3. Peak tides **Victoria:** November 15-17, December 12-14, December 29-31, January 8-11 and February 8.
4. Peak tides **Vancouver:** November 17-19, December 15-19 and January 12-16.
5. **SSH anomalies:** Near zero on the coast but positive in the north Pacific.

Mixed meteorological conditions are present early this fall already accompanied by several strong short-lived storms. Seasonal high tides will occur starting in mid-November for the Lower Mainland and Victoria, and be prominent in mid-December and mid-January for all areas of the south coast. La Niña and negative PDO conditions have worked together in the past to generate active storm seasons and may impact the strength and frequency of storms along the southern British Columbia coast this winter.

BC Storm Surge Forecasting System

References

- Eichler T. and W. Higgins, 2006: Climatology and ENSO-related variability of North American extratropical cyclone activity. *J. Clim* 19:2076-2093
- Gershunov, A. and T. P. Barnett. 1998. Interdecadal modulation of ENSO teleconnections. *Bulletin of the American Meteorological Society* 79: 2715-2725
- Mantua, N. J. and S. R. Hare. 2002. The Pacific Decadal Oscillation. *J. Ocean.* 58: 35-44
- Miller, A. J., D. R. Cayan, T. P. Barnett, N. E. Graham and J. M. Oberhuber, 1994. The 1976-77 climate shift of the Pacific Ocean. *Oceanography*, 7, 21-26
- Weng, H., Behera, S. K., & Yamagata, T. , 2009. Anomalous winter climate conditions in the Pacific rim during recent El Niño Modoki and El Niño events. *Climate Dynamics*, 32(5), 663-674.
- Wolter, K., and M. S. Timlin, 2011: El Niño/Southern Oscillation behaviour since 1871 as diagnosed in an extended multivariate ENSO index (MEI.ext). *Intl. J. Climatology*, 31, 14pp., in press.

BC Storm Surge Forecasting System

Appendix

Pacific Decadal Oscillation

The PDO, which represents the principal mode of variability of sea surface temperature in the Pacific Ocean, shifts between a positive (warm) and negative (cool) phase (Mantua and Hare, 2002). During a warm PDO phase, the waters off the west coast of North America exhibit warmer than normal temperatures, and wind velocity anomalies along the outer coast are directed northward (Fig. A1). The opposite is true during the cool phase. The time series of PDO index from 1900 to present (Fig. A2) shows that the oscillation between the warm and cool phase varies with a period of roughly 5-10 years, but can also stay predominantly in one mode for prolonged periods. The shift from a nearly 30-year cool phase to a warm phase in the late 1970s is considered by many to be an oceanic regime shift affecting both weather and biological ecosystems in the Pacific Ocean. After 1998, there was a return to the cool PDO phase. **NOTE: the JISAO PDO index is no longer updating as of September 2018; continuing data will be derived from the Japan Meteorological Agency which publishes PDO values derived from a parallel method to the Mantua index.**

The sea level pressure anomaly associated with the warm phase of the PDO acts to intensify the Aleutian low, and vice-versa during the cold phase. Gershunov and Barnett (1998) provide evidence of modulation of the ENSO signal by the PDO. The climate impacts of El Niño may be intensified during a warm PDO phase since both contribute to a deepening of the Aleutian low pressure system. Conversely, a cold phase PDO could enhance the effects of La Niña over western North America.




Figure A1. PDO warm and cool phases of sea surface temperature anomaly and associated wind anomaly patterns. Web image from Joint Institute for the Study of the Atmosphere and Ocean (JISAO), University of Washington.

BC Storm Surge Forecasting System




Figure A2. Monthly values of the PDO index from 1910-2020.

El Niño/La Niña

The El Niño-Southern Oscillation is a coupled tropical atmospheric and ocean phenomenon that has extra-tropical influence over weather and oceanographic conditions. The Southern Oscillation is a quasi-periodic fluctuation in the atmospheric sea level pressure measured between Tahiti (central south Pacific) and Darwin, Australia. When atmospheric pressure is lower than normal over Tahiti and higher than normal at Darwin, the Southern Oscillation Index (departures from the mean difference) is negative and the normally easterly trade winds are reduced. This causes the warm water mass normally pushed to the western side of the Pacific Ocean to migrate towards South America. The occurrence of warm water off the coast of Peru is known as El Niño, and coincides with abnormally high coastal sea levels which suppress coastal upwelling and adversely affect marine biota through reduced deep water nutrient supply. Extra-tropical effects of El Niño include warmer ocean and air temperatures on the west coast of Canada, and more southern storm tracks. The opposite effect (a strengthening of the trade winds) results in La Niña conditions, which can result in colder air, more precipitation and a northward shift of the Pacific storm track.

The impacts of El Niño/La Niña on sea levels off the coast of British Columbia can be significant. During the major El Niño years of 1982-83 and 1997-98, coastal sea level anomalies of 10-20 cm persisted for several months and contributed to a high occurrence of potentially damaging surges. La Niña can have the opposite effect (negative anomalies) but the anomalies are often more localized and of shorter duration. Figure A3 shows two images of residual sea surface height under both conditions.

BC Storm Surge Forecasting System




Figure A3. Residual sea surface height during the El Niño winter of 1997 and La Niña winter of 1999 (courtesy Jet Propulsion Laboratory). The positive SSH residual off the BC coast in 1997 is part of a larger feature stretching to the equator, while the negative SSH residual in 1999 is more localized. The magnitude of both is well in excess of 10 cm.

The Earth System Research Laboratory (ERSL) of the US National Oceanic and Atmospheric Administration produces a multi-variate ENSO index based on six oceanic and atmospheric variables: sea surface temperature, wind velocity components, sea level pressure, surface air temperature and cloudiness (Wolter and Timlin, 2011). The result is a monthly time series showing the interannual variability of ENSO strength (Fig. A4).

BC Storm Surge Forecasting System




Figure A4. NOAA ERSI multivariate ENSO (version 2) index from 1960 to present. Positive values indicate El Niño conditions, negative values indicate La Niña conditions.